



the art and science of smart patch testing™

NA68 Desoximetasone Patient Information

Your patch test result indicates that you have a contact allergy to desoximetasone. This contact allergy may cause your skin to react when it is exposed to this substance although it may take several days for the symptoms to appear. Typical symptoms include redness, swelling, itching and fluid-filled blisters.

Where is desoximetasone found?

Desoximetasone is a Class 3 high potency topical steroid that is available in an ointment or a gel. Desoximetasone is used to treat redness, swelling, itching, and discomfort of various skin conditions. Desoximetasone works by activating natural substances in the skin to reduce swelling, redness, and itching

How can you avoid contact with desoximetasone?

Avoid products that list any of the following names in the ingredients:

- | | |
|--|--------------------------------|
| ■ 17-Deoxymethasone | ■ Desoximetasone |
| ■ 17-Desoximethasone | ■ Desoximetasonium [INN-Latin] |
| ■ 9-Fluoro-11-beta,21-dihydroxy-16-alpha-methylpregna-1,4-diene-3,20-dione | ■ Desoxymethasone |
| ■ 9-Fluoro-11beta,21-dihydroxy-16alpha-methylpregna-1,4-diene-3,20-dione | ■ EINECS 206-845-3 |
| ■ A 41-304 | ■ Esperson |
| ■ A 41304 | ■ Flubason |
| ■ BRN 2228097 | ■ HOE 304 |
| ■ Deoxymethasone | ■ Ibaril |
| ■ Desossimetasone [DCIT] | ■ R 2113 |
| ■ Desoximetasona [INN-Spanish] | ■ Stiedex |
| | ■ Topicort |
| | ■ Topisolon |

What are some products that may contain desoximetasone?

Topical Medications:

- Generic desoximetasone ointments
- Topicort
- Topicort LP

*For additional information about products that might contain **desoximetasone**, go to the Household Product Database online (<http://householdproducts.nlm.nih.gov>) at the United States National Library of Medicine. These lists are brief and provide just a few examples. They are not comprehensive. Product formulations also change frequently. Read product labels carefully and talk to your doctor if you have any questions. These are general guidelines. Talk to your doctor for more specific instructions.